

Technical Note

Arthroscopic Treatment of the Arthrofibrotic Knee

David H. Kim, M.D., Thomas J. Gill, M.D., and Peter J. Millett, M.D., M.Sc.

Abstract: The management of motion loss of the knee is challenging. A clear understanding of the pathoanatomic causes of motion loss is necessary to establish a careful and rational approach to treatment. Early recognition and physical therapy are effective for the majority of patients, but when these conservative measures fail, operative intervention is indicated. The purpose of this article is to outline a comprehensive approach to the arthroscopic evaluation and treatment of the arthrofibrotic knee. This technique is designed to allow the surgeon to systematically address the numerous causes of motion loss of the knee. **Key Words:** Knee—Arthrofibrosis—Arthroscopy—Stiffness—Motion loss—Lysis of adhesions.

Motion loss of the knee can result from a variety of causes and poses a difficult clinical problem. Early recognition and appropriate treatment can be expected to restore motion and improve function in the majority of patients. An understanding of the pathoanatomic causes of motion loss is necessary to provide successful treatment strategies. Whenever possible, it is important to identify the specific cause and target treatment accordingly.

Nonoperative measures such as rest, ice, anti-inflammatory agents, and subsequent physical therapy are sometimes unsuccessful. In such settings, operative intervention is indicated. Manipulation under anesthesia has been used in the postoperative period to address motion loss with some success.^{1,2} However, manual manipulation has significant risk, including complications such as distal femoral fracture and patellar tendon rupture, but this can be prevented by a

controlled surgical release of scar tissue as opposed to a gross manual manipulation under anesthesia. The arthroscopic approach is a powerful and controlled method that is effective both for focal, discrete lesions as well as for more global arthrofibrosis. The purpose of this article is to outline our systematic approach to the arthroscopic surgical management of the arthrofibrotic knee.

GENERAL PRINCIPLES

Fibrosis and contractures in different parts of the knee contribute to different types of motion loss. Adhesions in the suprapatellar pouch typically limit patellar mobility and can restrict knee flexion. The proximal extent of the pouch should be approximately 3.5 cm from the superior pole of the patella. A foreshortened pouch can lead to a further loss of knee flexion.

Other structures of the knee that contribute to a loss of flexion are the medial and lateral gutters, and the anterior interval. The anterior interval is the region of the knee posterior to the patellar fat pad and anterior to the anterosuperior tibial plateau. This interval is an underrecognized source of knee flexion loss.

The cruciate ligaments themselves can lead to restrictions in motion, particularly after cruciate ligament reconstruction.³⁻⁸ An anteriorly placed femoral tunnel^{9,10} or overtensioned graft^{11,12} can limit knee

From the Department of Orthopaedic Surgery (D.H.K., P.J.M.), Brigham and Women's Hospital, Boston, Massachusetts; and the Department of Orthopaedic Surgery (D.H.K., T.J.G., P.J.M.), Massachusetts General Hospital, Boston, Massachusetts, U.S.A.

Address correspondence and reprint requests to Peter J. Millett, M.D., M.Sc., Brigham and Women's Hospital, 75 Francis Street, Boston, MA 02115, U.S.A. E-mail: pmillett@partners.org

*© 2004 by the Arthroscopy Association of North America
0749-8063/04/2006-3913\$30.00/0*

doi:10.1016/j.arthro.2004.04.036




FIGURE 1. Coronal (A) and sagittal (B) illustrations of the knee. Schematic diagram of nine-step systematic arthroscopic evaluation: (1) evaluate and re-establish the suprapatellar pouch; (2) evaluate and re-establish the medial gutter; (3) evaluate and re-establish the lateral gutter; (4) mobilize and debride the infrapatellar fat pad to re-establish the pretibial recess; (5) evaluate the lateral retinaculum and perform lateral retinacular release, if necessary; (6) evaluate the medial retinaculum and perform medial retinacular release, if necessary; (7) evaluate the intracondylar notch, debride scar tissue, and in severe cases, release the anterior cruciate ligament/posterior cruciate ligament, if necessary; (8) evaluate the tibial insertion of posterior capsule, inspect the capsular recess, and perform medial and lateral capsulotomy, if necessary; and (9) evaluate the femoral insertion of the posterior capsule and release capsule, if necessary.

flexion. An anteriorly placed anterior cruciate ligament (ACL) graft on the tibia will impinge on the intercondylar notch and limit knee extension. This impingement could lead to the formation of fibrous ACL nodules (also known as “cyclops” lesions⁴), which further limit extension and can worsen impingement.^{4,13-17}

A tight posterior knee capsule can limit knee extension. This area is difficult to approach arthroscopically and could be best addressed through an open approach if needed.

When surgery is undertaken, we perform a systematic nine-step evaluation¹⁸ (Fig 1). First, adhesions in the suprapatellar pouch are identified and released. We seek to restore the normal suprapatellar pouch to allow excursion between the extensor mechanism and the underlying tissues. Next, the medial and lateral gutters are inspected and releases are performed as needed. The anterior interval should be evaluated dynamically as the knee is placed through flexion and extension. This interval between the patellar fat pad and anterior tibia is released to re-establish the pretibial recess. Next, we evaluate the lateral and

medial retinacular structures and perform selective releases of any tight lateral and medial retinacular bands of tissue. A complete retinacular release is needed only in severe cases. The intercondylar notch is debrided of scar tissue, particularly fibrous ACL nodules, which lead to notch impingement. In overt cases of graft malposition, the ACL or even the posterior cruciate ligament (PCL) might need to be released. Knee motion is then assessed. If full extension cannot be achieved, the tibial insertion of the posterior capsule is inspected. If the posterior capsule is tight, medial, and occasionally lateral capsulotomies are performed. Finally, we evaluate the femoral insertion of the posterior capsule if full extension is not achieved and an open posteromedial release is performed if needed.

OPERATIVE TECHNIQUE

Anesthesia

We recommend the use of regional epidural anesthesia and the placement of an indwelling epidural



FIGURE 2. Capsular distention: Using a 60-mL syringe and an 18-gauge needle, a total of 180 mL of normal saline is introduced into the knee joint before the placement of the arthroscopic portals.

catheter for intraoperative and postoperative analgesia. This type of anesthesia provides improved localized pain control and allows more intensive physical therapy in the immediate postoperative period.

Capsular Distention

Capsular distention with 120 to 180 mL of saline before arthroscopy is a useful adjunct in the arthroscopic treatment of arthrofibrosis of the knee.¹⁹ With the patient under anesthesia and after the extremity is prepped and draped, the knee joint is palpated. Often the scarring is so severe that it is difficult to make out the usual surface landmarks. Using an 18-gauge needle and 60-mL syringe, normal saline is injected into the suprapatellar pouch from the lateral side (Fig 2). It is important to carefully watch and feel for joint distention as the fluid is injected to ensure that the fluid is entering the true joint space. Fluid should flow easily if the needle is indeed in the true joint space. Normal knees easily accept 180 mL of saline, and an attempt to introduce this volume of fluid into the arthrofibrotic knee should be made. As the capsule is distended with fluid, care is taken to avoid rupturing the true capsule, although intra-articular adhesions could be disrupted. The last 60 mL of fluid is inserted slowly to allow the capsule to stretch over time. Preservation of the true joint capsule prevents extravasa-

tion of the fluid during arthroscopy and facilitates visualization.

Portal Placement/Visualization

After the knee is maximally distended with saline, we typically insert an inflow cannula into the knee through a superolateral portal and then initiate flow. This keeps the joint distended and facilitates insertion of the arthroscope through the standard inferolateral portal and also helps with initial visualization of the joint. In severe cases, it is often difficult simply to insert the arthroscope. A standard inferomedial working portal is then established. Although the camera sheath can be used to manually release adhesions, we prefer to use electrocautery to lyse adhesions. This approach will minimize bleeding, which not only helps with visualization, but also decreases the risk of recurrent postoperative scarring.

Systematic Examination

Suprapatellar Pouch: Using electrocautery, we lyse adhesions and release scarring to re-establish the suprapatellar pouch (Fig 3). It is important to remember that the pouch is quite large and should extend 3 to 4 cm proximal to the patella and releases should continue until this is achieved.

Medial and Lateral Gutters: It is important to look for adhesions in this region. Often one can see the adhesions that have formed between the capsule and the femoral condyles. A suction punch or electrocautery can be used to remove adhesions from the lateral gutter.

Anterior Interval: The infrapatellar fat pad and pretibial recess are inspected. We release and mobilize the infrapatellar fat pad from the anterior tibia and re-establish the pretibial recess (Fig 4). With a normal anterior interval, it is important to notice how the intermeniscal ligament glides freely on the anterior surface of the tibia. With anterior interval scarring, adhesions form in the pretibial recess. In experimental models,²⁰ these adhesions have been shown to increase patellofemoral joint contact pressures. In such instances, an anterior interval release should be performed by releasing the scar tissue from medial to lateral just anterior to the peripheral rim of the anterior horn of each meniscus. We prefer to use bipolar tissue ablation. The release also proceeds from the proximal extent at the level of the meniscus to approximately 1 cm distal along the anterior tibial cortex. Care should be taken to avoid cauterizing or burning the bone of the anterior tibia. Meticulous hemostasis should be maintained, especially around the infrapatellar fat pad.



FIGURE 3. Re-establishment of the suprapatellar pouch: Adhesions and scarring are lysed using electrocautery to re-establish the suprapatellar pouch.

Lateral and Medial Retinaculum: Using electrocautery or a suction punch, we perform selective releases of lateral and medial retinacular structures if they are tight or scarred. This decompresses the patellofemoral joint and improves patellofemoral mobility. It also enhances visualization and permits easier instrumentation as the effective joint space is increased.

Intercondylar Notch: If there is evidence of graft impingement, a notchplasty should be performed. Fibroproliferative ACL nodules (Fig 5A) should be excised using a suction punch, because these nodules can limit extension by impinging on the intercondylar notch. If the cruciate grafts are malpositioned or the native ligaments are excessively scarred, they can be

FIGURE 4. Re-establishment of the anterior interval: The infrapatellar fat pad and pretibial recess are debrided using an electrocautery device.






FIGURE 5. (A) Fibroproliferative anterior cruciate ligament nodule: Impingement can be caused by the cyclops lesion. (B) Bony nodule: An arthroscopic bur can be used to remove a bony nodule on the anterior cruciate ligament.

debrided, released, or excised altogether, according to the severity of involvement. Often patients will present with a bony nodule in the intercondylar notch. Such nodules or osteophytes can also impinge in the intercondylar notch and will block knee extension. A high-speed arthroscopic bur (Fig 5B) or an osteotome can be used to remove these bony nodules to prevent further impingement and to re-establish full extension. Generally, peripheral osteophytes are not debrided because this causes unnecessary bleeding and potentiates further adhesions.

Posterior Capsule: After completing the evaluation of the intercondylar notch, the knee should be placed through a range of motion, and flexion and extension should be carefully assessed. If there is a persistent loss of extension, consideration should be made for an open capsular release. We prefer to perform this using a limited open approach as described by Steadman et al.²¹ Such releases to the posterior medial and posterior lateral capsules should be added if the posterior capsule remains tight. In addition to a capsulotomy, a blunt periosteal elevator is used to

strip the capsule from the tibial and femoral insertions. Generally, we begin on the medial side and release the capsule here. If a loss of extension persists, a similar open lateral capsular release is performed.

In our experience, postoperative knee motion will not improve from that obtained in surgery. Therefore, if a loss of extension or flexion persists intraoperatively, every attempt should be made to regain this motion before leaving the operating room.

Postoperative Protocol/Rehabilitation

Patients are typically hospitalized for 48 hours to take advantage of the analgesic properties provided by the indwelling epidural catheter. In the immediate postoperative period, patients are placed in a continuous passive motion (CPM) machine for at least 8 to 10 hours a day to maintain the range of motion obtained in the operating room. CPM is less effective than manual mobilization for the terminal ranges of motion. Home CPM is continued for 2 to 3 weeks. Dynamic bracing can be used to maintain full extension, although these devices are occasionally not well tolerated. Patellar mobility exercises, extensor mechanism exercises, and full passive and active-assisted range of motion exercises are all essential components of therapy, which begin on the first postoperative day.

Daily outpatient physical therapy visits start when patients are discharged and could continue for 6 to 8 weeks. Stationary bicycling could begin immediately, whereas aquatherapy and resistance exercises are begun at 2 weeks if range of motion is maintained and swelling is controlled.

DISCUSSION

The term arthrofibrosis is often used to describe loss of flexion, loss of extension, or both. More precisely, it refers to a specific process in which scar tissue or fibrous adhesions form diffusely within a joint.²² Risk factors to developing arthrofibrosis of the knee include multiligament injuries and knee dislocations,²³⁻²⁷ as well as prolonged immobilization, infection, and reflex sympathetic dystrophy.^{28,29}

It is important to understand the myriad causes of motion loss so that treatment can be targeted at the specific cause. Causes of extension loss include notch impingement, ACL nodules, and posterior capsule pathology. Causes of flexion loss include suprapatellar adhesions, as well as medial and lateral gutter adhesions. Causes of both extension and flexion loss include improper graft position, infrapatellar contracture

syndrome,³⁰⁻³² soft tissue calcifications, global arthrofibrosis, reflex sympathetic dystrophy, and infection. As a general rule, causes of loss of extension reside in the intercondylar notch and posterior capsule, whereas causes of loss of flexion reside in the suprapatellar pouch and the medial and lateral gutters.

The arthroscopic treatment of knee arthrofibrosis has been reported as being effective in improving knee range of motion and restoring function with minimal complications. Among the first to report their results were Sprague et al.,³³ who described a series of 24 patients with "fibroarthrosis" who gained an average of 45° of knee flexion with arthroscopic release of adhesions. Several authors have since reported similar results with knee range-of-motion improvements ranging from 45° to 68°.^{29,33-42}

In addition, several reports have reviewed the results of arthroscopic treatment of arthrofibrosis, specifically after ACL reconstruction.^{4,8,14,15,43} Jackson and Schaefer⁴ first described the "cyclops" lesion after ACL reconstruction and reported on 13 patients who underwent arthroscopic debridement with knee manipulation and achieved an average improvement of loss of extension from 16° to 3.8°. Marzo et al.¹⁴ reported on 21 patients who improved on an average loss of extension of 11° to an average of 0° at 1-year follow up after arthroscopic removal of fibrous nodules in the intercondylar notch. Subsequent studies^{8,15,43} have all demonstrated marked improvements in function and symptoms with arthroscopic debridement of anterior scar formation in combination with aggressive rehabilitation and serial extension casting.

Arthroscopic treatment is generally most effective when the underlying pathology is predominately intra-articular. When the etiology of the motion loss is multifactorial or involves extra-articular structures, often a combined approach involving both arthroscopic and open techniques is indicated.^{21,44-46} Results from this type of treatment are generally less satisfactory than with those reported from solely an arthroscopic approach, mainly because the underlying causes for motion loss are more extensive.

SUMMARY

Arthrofibrosis has a variety of causes and for optimal outcomes, appropriate treatment should be targeted at the specific cause. Our nine-step arthroscopic approach allows the surgeon to systematically address all of the intra-articular pathoanatomy. The first three steps are to evaluate and re-establish the suprapatellar pouch followed by the medial and lateral gutters.

Next, the anterior interval, infrapatellar fat pad, and pretibial recess should be re-established. The medial and lateral retinaculum are evaluated and releases performed if necessary. Electrocautery is essential to minimize bleeding. The intercondylar notch must be carefully evaluated, because this is often a major site of pathology. Scar tissue, ACL nodules, bony nodules, or malpositioned grafts can all cause motion problems. The final two steps involve evaluation of the medial and lateral posterior capsule at both the tibial and femoral insertions. For those patients with persistent loss of extension after comprehensive arthroscopic treatment, limited open releases should be undertaken.

REFERENCES

- Dodds JA, Keene JS, Graf BK, et al. Results of knee manipulations after anterior cruciate ligament reconstructions. *Am J Sports Med* 1991;19:283-287.
- Noyes FR, Mangine RE, Barber SD. The early treatment of motion complications after reconstruction of the anterior cruciate ligament. *Clin Orthop* 1992;277:217-228.
- Johnson RJ, Eriksson E, Haggmark T, et al. Five- to ten-year follow-up evaluation after reconstruction of the anterior cruciate ligament. *Clin Orthop* 1984;183:122-140.
- Jackson DW, Schaefer RK. Cyclops syndrome: Loss of extension following intra-articular anterior cruciate ligament reconstruction. *Arthroscopy* 1990;6:171-178.
- Shelbourne KD, Wilckens JH, Mollabashy A, et al. Arthrofibrosis in acute anterior cruciate ligament reconstruction. The effect of timing of reconstruction and rehabilitation. *Am J Sports Med* 1991;19:332-336.
- Harner CD, Irrgang JJ, Paul J, et al. Loss of motion after anterior cruciate ligament reconstruction. *Am J Sports Med* 1992;20:499-506.
- Cosgarea AJ, Sebastianelli WJ, DeHaven KE. Prevention of arthrofibrosis after anterior cruciate ligament reconstruction using the central third patellar tendon autograft. *Am J Sports Med* 1995;23:87-92.
- Shelbourne KD, Patel DV, Martini DJ. Classification and management of arthrofibrosis of the knee after anterior cruciate ligament reconstruction. *Am J Sports Med* 1996;24:857-862.
- Romano VM, Graf BK, Keene JS, et al. Anterior cruciate ligament reconstruction. The effect of tibial tunnel placement on range of motion. *Am J Sports Med* 1993;21:415-418.
- Fu FH, Bennett CH, Lattermann C, et al. Current trends in anterior cruciate ligament reconstruction. Part I: Biology and biomechanics of reconstruction. *Am J Sports Med* 1999;27:821-830.
- Melby AD, Noble JS, Askew MJ, et al. The effects of graft tensioning on the laxity and kinematics of the anterior cruciate ligament reconstructed knee. *Arthroscopy* 1991;7:257-266.
- Nabors ED, Richmond JC, Vannah WM, et al. Anterior cruciate ligament graft tensioning in full extension. *Am J Sports Med* 1995;23:488-492.
- Fullerton LR Jr, Andrews JR. Mechanical block to extension following augmentation of the anterior cruciate ligament. A case report. *Am J Sports Med* 1984;12:166-168.
- Marzo JM, Bowen MK, Warren RF, et al. Intraarticular fibrous nodule as a cause of loss of extension following anterior cruciate ligament reconstruction. *Arthroscopy* 1992;8:10-18.
- Fisher SE, Shelbourne KD. Arthroscopic treatment of symptomatic extension block complicating anterior cruciate ligament reconstruction. *Am J Sports Med* 1993;21:558-564.
- Recht MP, Piraino DW, Cohen MA, et al. Localized anterior arthrofibrosis (cyclops lesion) after reconstruction of the anterior cruciate ligament: MR imaging findings. *AJR Am J Roentgenol* 1995;165:383-385.
- Delince P, Krallis P, Descamps PY, et al. Different aspects of the cyclops lesion following anterior cruciate ligament reconstruction: A multifactorial etiopathogenesis. *Arthroscopy* 1998;14:869-876.
- Millett PJ, Wickiewicz TL, Warren RF. Motion loss after ligament injuries to the knee. Part II: Prevention and treatment. *Am J Sports Med* 2001;29:822-828.
- Millett PJ, Steadman JR. The role of capsular distention in the arthroscopic management of arthrofibrosis of the knee: A technical consideration. *Arthroscopy* 2001;17:E31.
- Ahmad CS, Kwak SD, Ateshian GA, et al. Effects of patellar tendon adhesion to the anterior tibia on knee mechanics. *Am J Sports Med* 1998;26:715-724.
- Steadman JR, Burns TP, Pelozo J. Surgical treatment of arthrofibrosis of the knee. *J Orthop Tech* 1993;1:119-127.
- Petsche TS, Hutchinson MR. Loss of extension after reconstruction of the anterior cruciate ligament. *J Am Acad Orthop Surg* 1999;7:119-127.
- Sisto DJ, Warren RF. Complete knee dislocation. A follow-up study of operative treatment. *Clin Orthop* 1985;198:94-101.
- Ekeland A, Vikne J. Treatment of acute combined knee instabilities and subsequent sport performance. *Knee Surg Sports Trauma Arthrosc* 1995;3:180-183.
- Shapiro MS, Freedman EL. Allograft reconstruction of the anterior and posterior cruciate ligaments after traumatic knee dislocation. *Am J Sports Med* 1995;23:580-587.
- Noyes FR, Barber-Westin SD. Reconstruction of the anterior and posterior cruciate ligaments after knee dislocation. Use of early protected postoperative motion to decrease arthrofibrosis. *Am J Sports Med* 1997;25:769-778.
- Cole BJ, Harner CD. The multiple ligament injured knee. *Clin Sports Med* 1999;18:241-262.
- O'Brien SJ, Ngeow J, Gibney MA, et al. Reflex sympathetic dystrophy of the knee. Causes, diagnosis, and treatment. *Am J Sports Med* 1995;23:655-659.
- Lindenfeld TN, Wojtys EM, Husain A. Instructional Course Lectures, the American Academy of Orthopaedic Surgeons—operative treatment of arthrofibrosis of the knee. *J Bone Joint Surg Am* 1999;81:1772-1784.
- Paulos LE, Rosenberg TD, Drawbert J, et al. Infrapatellar contracture syndrome. *Am J Sports Med* 1987;15:331-341.
- Noyes FR, Wojtys EM, Marshall MT. The early diagnosis and treatment of developmental patella infera syndrome. *Clin Orthop* 1991;265:241-252.
- Paulos LE, Wnorowski DC, Greenwald AE. Infrapatellar contracture syndrome. Diagnosis, treatment, and long-term followup. *Am J Sports Med* 1994;22:440-449.
- Sprague NF III, O'Connor RL, Fox JM. Arthroscopic treatment of postoperative knee arthrofibrosis. *Clin Orthop* 1982;166:165-172.
- DelPizzo W, Fox JM, Friedman MJ, et al. Operative arthroscopy for the treatment of arthrofibrosis of the knee. *Contemporary Orthopaedics* 1985;10:67-72.
- Sprague NF III. Motion-limiting arthrofibrosis of the knee: the role of arthroscopic management. *Clin Sports Med* 1987;6:537-549.
- Parisien JS. The role of arthroscopy in the treatment of postoperative fibroarthrosis of the knee joint. *Clin Orthop* 1988;229:185-192.
- Richmond JC, al Assal M. Arthroscopic management of arthrofibrosis of the knee, including infrapatellar contraction syndrome. *Arthroscopy* 1991;7:144-147.

38. Achalandabaso J, Albillos J. Stiffness of the knee—mixed arthroscopic and subcutaneous technique: results of 67 cases. *Arthroscopy* 1993;9:685-690.
39. Cohen I, Hendel D, Rzetelny V. Arthroscopic adhesiolysis of the knee joint in arthrofibrosis. *Bull Hosp Jt Dis* 1993;53:66-67.
40. Vaquero J, Vidal C, Medina E, et al. Arthroscopic lysis in knee arthrofibrosis. *Arthroscopy* 1993;9:691-694.
41. Cosgarea AJ, DeHaven KE, Lovelock JE. The surgical treatment of arthrofibrosis of the knee. *Am J Sports Med* 1994;22:184-191.
42. Klein W, Shah N, Gassen A. Arthroscopic management of postoperative arthrofibrosis of the knee joint: Indication, technique, and results. *Arthroscopy* 1994;10:591-597.
43. Shelbourne KD, Johnson GE. Outpatient surgical management of arthrofibrosis after anterior cruciate ligament surgery. *Am J Sports Med* 1994;22:192-197.
44. Aglietti P, Buzzi R, De Felice R, Paolini G, Zaccherotti G. Results of surgical treatment of arthrofibrosis after ACL reconstruction. *Knee Surg Sports Trauma Arthrosc* 1995;3:83-88.
45. Lobenhoffer HP, Bosch U, Gerich TG. Role of posterior capsulotomy for the treatment of extension deficits of the knee. *Knee Surg Sports Trauma Arthrosc* 1996;4:237-241.
46. Millett PJ, Williams RJ III, Wickiewicz TL. Open debridement and soft tissue release as a salvage procedure for the severely arthrofibrotic knee. *Am J Sports Med* 1999;27:552-561.